



INTERNATIONAL JOURNAL OF PURE AND APPLIED RESEARCH IN ENGINEERING AND TECHNOLOGY

A PATH FOR HORIZING YOUR INNOVATIVE WORK

DEGRADATION OF M-CHLOROPHENOL USING SOLAR RADIATIONS

MRS. SHILPA S. PATIL¹, PROF. S. T. PATIL², MR. ROHIT SHETTY³

1. M.TECH Student (Structure and Construction) Ballarpur Institute of Technology, Ballarpur.
2. M.TECH Student (Structure and Construction) Ballarpur Institute of Technology, Ballarpur.
3. M.TECH Student (Structure and Construction) Ballarpur Institute of Technology, Ballarpur.

Accepted Date: 05/03/2015; Published Date: 01/05/2015

Abstract: The degradation of m-chlorophenol of was investigated in presence of commercially available Degussa P-25TiO₂ photocatalyst and solar radiation as well as artificial UV irradiation. The objective of present article is to study the advanced separation techniques of m-chlorophenol present in pharmaceutical wastewater and to avoid the future accumulation of m-chlorophenol in aquatic environment. The effect of different operating parameters like initial concentration of m-chlorophenol catalytic loading, pH of solution, effect of anion and captions on PCD of m-chlorophenol have been investigated in detail. It was observed that as initial concentration of m-chlorophenol increase rate of PCD decreases. Optimum catalytic loading was observed at 1 g/L for m-chlorophenol concentration 0.1gm/l. From kinetic studies, it was found that the PCD obeys pseudo-first order kinetics with reaction rate constant.

Keywords: Photocatalysist, Solar, Artificial Radiations.

Corresponding Author: MRS. SHILPA S. PATIL



PAPER-QR CODE

Access Online On:

www.ijpret.com

How to Cite This Article:

Shilpa S. Patil, IJPRET, 2015; Volume 3 (9): 21-29

INTRODUCTION

The rapid growth of the world's population has created an increase in the demand for the Earth's limited supply of freshwater. Thus, protecting the supply of water and addressing concerns such as toxicity and the presence of chemicals that may have potential long term adverse human and ecological effects has become an important issue. Pharmaceuticals and personal care products enter the environment at low concentrations primarily as metabolites excreted by humans and animals or in effluents that are disposed into wastewater from hospitals, pharmacies, and chemical manufacturing facilities. Once these compounds reach wastewater treatment plants, they are not completely removed and presence of m-chlorophenol in aquatic environments may cause serious threats to the ecosystem and human health [3, 4].

Therefore there is a need to develop eco-friendly, cost effective technique for the treatment of pharmaceutical wastewater. The studies have shown that advanced oxidation processes (AOP) and adsorption are viable treatment methods for completely removing m-chlorophenol from water and wastewater. Recently, extensive studies have been performed on the degradation of toxic organic compounds in waste water via modified semiconductor photocatalyst [5]. The purpose of this article is to investigate the photocatalytic degradation (PCD) of m-chlorophenol using Degussa P-25 TiO_2 photocatalyst and using solar and artificial radiation. The effect of various operating parameters such as initial concentration of m-chlorophenol, photocatalyst loading, pH of solution and effect of presence of co-existing ion on photocatalytic oxidation of m-chlorophenol have been studied in order to establish the optimum process parameters. The kinetics of photocatalytic oxidation has been also studied

2. MATERIAL AND METHODS

2.1 Reagent -Aeroxide Degussa P-25 TiO_2 (70:30% w/w anatase to rutile) with an average particle size of 30 nm and BET surface area about $55 \text{ m}^2 \text{ g}^{-1}$ was used as the photocatalyst. Acetic acid (CH_3COOH), Acetonitrile, Na_2CO_3 , NaCl and $(\text{NH}_4)_2\text{SO}_4$ were purchased from Merck India Pvt. Ltd, Mumbai, India. The different concentration aqueous solution of m-chlorophenol was prepared in de-ionized water (DI Water) was used as pollutant.

2.2 Equipment

A parabolic reflector with surface area of 1.5 m^2 was used for concentration of solar radiation. The pH of aqueous solution of ciprofloxacin HCl was monitored by using pH meter of Thermo scientific Orion star model. Plain solar radiation intensity was measured in W m^{-2} by 'daystar meter' (daystar Inc, Las Cruces, NM, USA) working on the photocell principle. Solar

radiation intensity at ground level is referred to as plain intensity henceforth. Average solar intensity of solar irradiation was found approximately 800 watts/m². Experimental setup is as shown in Fig no.2. Deionized water (Millipore Milli-Q system) was used to prepare aqueous solutions of ciprofloxacin HCl. Cooling media (water) was circulated by chiller (Julabo).

2.3 PCD Experiment

An annular borosilicate glass reactor as shown in Figure 2 was used as photocatalytic for performing the photocatalytic oxidation reaction. The de-ionized water (Millipore Milli-Q system) was used to prepare aqueous solutions of m-chlorophenol. The temperature of reaction mixture was kept constant by circulated the chilled water through the annular space between the lamp via JULABO chiller FP-50 MA. In each experiment 500 ml of solution was charged in the reactor. An air at rate of 1 L min⁻¹ required for oxidation reaction and to keep all the TiO₂ in suspension was supplied via ring sparger located at the bottom of the reactor. In the photocatalytic experiments Degussa P-25 TiO₂ was used as the photocatalyst. The solution was equilibrated by stirring for 15 minutes in the dark before exposing the reactor assembly to the solar radiation. The liquid samples were taken periodically from the reactor for analysis and stored in amber colored bottle. During the PCD experiments the pH of the solution was not controlled. Same experimental procedure was carried out by using artificial radiations. A visible tungsten lamp with power input 400 watts was used as the source of artificial radiations. shown in fig.1




Fig 1. Photocatalytic reactor for Artificial method



Figure 2: Experimental Setup

3. RESULTS AND DISCUSSION

3.1 Initial concentration effect

Initial concentration of pollutant is the important parameter that affects the rate of PCD of pollutant. The experiments on PCD of meta-chlorophenol were conducted for solutions with range of 0.1gm/l – 1 gm/l concentrations. It was found that meta-chlorophenol of initial concentration 50 mg/l was completely degraded within 2 hr by using solar radiations. The PCD time increased with increase in meta-chlorophenol concentration. In this particular study catalyst concentration was kept constant 1 gm/l. The concentration of meta-chlorophenol has a significant effect on the PCD. The rate was higher when the initial concentration meta-chlorophenol was less. This can be explained as for a certain TiO_2 concentration, the amount of active centers on the photocatalyst was finite, so PCD rate was fast with low concentration of meta-chlorophenol. The PCD rate decreases when the concentration of increases, possibly because the molecules of meta-chlorophenol was excessive in comparison with the amount of active centers on the photocatalyst. Figure shows the effect of initial concentration of meta-chlorophenol on rate of PCD. The photo catalytic process was influenced by the initial concentration of meta-chlorophenol A linear relationship was observed between meta-chlorophenol concentration and irradiation time, as shown in Figure 2.




Figure 2: Effect of initial concentrations

3.2 Effect of catalyst concentration on rate of degradation-The concentration of TiO_2 is a critical parameter in photo catalytic oxidation. It effects on the reaction rate and consequently on the cost of treatment. The effect of catalyst loading of photocatalyst was studied on 0.1 gm/l of aqueous solution of ciprofloxacin HCl. The range of catalyst loading was 0 to 2 gm/l. It was observed that the pH of solution was 5.8 during this particular.




Figure 3: Effect of catalytic TiO₂

3.3 Effect of pH of solution on degradation of meta-chlorophenol -It was observed that PCD of m-chlorophenol was faster in weak basic (pH 9) than those in weak acidic (pH 5) or strong alkaline (pH 11) conditions [6]. It shows that the rate PCD of m-chlorophenol is strongly influenced by the reaction pH. The TiO₂ surface will remain positively charged in acidic medium (pH < 6.25) and negatively charged in alkaline medium (pH > 6.25) because of that reason under acidic condition (3 pH), m-chlorophenol is positively charged which is same as TiO₂ surface resulting in repulsion between them. Cl⁻ may compete with water molecules for the adsorption on TiO₂ surface this blocks the active sites of TiO₂ and hence production of hydroxyl radical decreases. Both TiO₂ and m-chlorophenol are negatively charged under strongly basic conditions (pH 11) which causes repulsion between them and lowers the PCD. Effect of pH on rate of degradation is as shown in Figure5. It was observed that at pH 9 rate of degradation was maximum.





Figure 4: Effect of pH on Rate

3.4 Effect of co-existing ions


The presence of certain inorganic ions in the reaction medium has a significant effect on degradation of pollutants. The effect of sodium carbonate, sodium chloride and ammonium sulphate on degradation m-chlorophenol was studied. During this experiment salt concentration was 0.5 gm/l. m-chlorophenol concentration was 0.1 gm/l. and 2 gm/l of TiO₂ was suspended for 120 min of radiation sodium carbonate show less degradation as compared to other salts. For 120 min run of salt addition experiment shows lower PCD rate of m-chlorophenol than that of solution without salt as shown in Figure 6.

For SO₄²⁻ ions, they are adsorbed in the surface of TiO₂ and consequently react with photo-induced holes (h⁺)



As 'S' is the strong oxidizing agent, the sulfate radical can accelerate the reaction and participate in the degradation.

For NaCl it reacts with m-chlorophenol to form other side product or it improves the adsorption of m-chlorophenol on surface of TiO₂. Na₂CO₃ has hindrance effect which reduces the m-chlorophenol adsorption onto the catalyst surface which results into the decreased PCD of m-chlorophenol.



4. KINETIC MODELS OF PHOTOCATALYTIC OXIDATION OF M-CHLOROPHENOL

Langmuir–Hinshelwood rate expression has been successfully used for heterogeneous photocatalytic degradation to determine the relationship between the initial degradation rate and the initial concentration of the organic substrate. The Langmuir–Hinshelwood model was developed to describe gas–liquid reaction it is widely used for liquid–solid reactions. Reaction

kinetics gives information about the reaction rates and the mechanisms by which the reactants are converted to the products.

PCD proceed on the surface of semiconductor via several steps. In this model, the reaction rate (r) is considered to be proportional to the surface fraction covered by the substrate (θ).


$$-\ln\left(\frac{C}{C_0}\right) = K_{app} * t$$

Radiation and solar radiation respectively. Regression coefficients are more than 0.96, it shows good fitting of data.

The effect of initial concentration of m-chlorophenol on PCD rate was studied by using solar and artificial radiation Table 1 shows the R^2 and K_{app} for artificial and solar radiation, from this Table it was found that K_{app} decreases as concentration of m-chlorophenol increases, but rate constant for solar radiation is much higher as compare to artificial radiation.

| Initial Conc. Of m-chlorophenol | Solar radiation | | Artificial radiation | |
|---------------------------------|-------------------|-------|----------------------|-------|
| | K_{app} (1/min) | R^2 | K_{app} (1/min) | R^2 |
| 25mg/l | 0.027 | 0.92 | - | - |
| 50mg/l | 0.026 | 0.95 | - | - |
| 75 mg/l | 0.022 | 0.90 | 0.020 | 0.82 |
| 100mg/l | 0.037 | 0.956 | 0.022 | 0.86 |

Table 1: Regression coefficients (R^2) and K_{app} for initial concentration



4. CONCLUSIONS

PCD of m-chlorophenol using solar and artificial radiation in the presence of TiO₂ photocatalyst was successfully accomplished. In this investigation it was found that K_{app} in the presence solar radiation is much higher than that of artificial radiation. This indicates that solar radiations are much more effective as compared to artificial radiations.

REFERENCES

1. M.Roma, M.Weller, S.Wentzell, Removal of Ciprofloxacin from Water using Adsorption, UV Photolysis and UV/H₂O₂ Degradation, Worcester Polytechnic Institute, BS Degree report April 28, 2011.
2. A. Nikolaou, S.Meri, D.Fatta, Occurrence patterns of pharmaceuticals in water and wastewater environments, Anal Bioanal Chem 387, 2007, 1225–1234.
3. T.Ternes, A.Joss, Human Pharmaceuticals, Hormones and Fragrances: The Challenge of Micropollutants in Urban Water Management, first ed., IWA, Cornwall, 2006.
4. A.Watkinson, E.Murby, S. Costanzo, Removal of antibiotics in conventional and advanced wastewater treatment: implications for environmental discharge and wastewater recycling, Water Res, 41, 2007, 4164–4176.
5. W.Zhao, W.Ma, C. Chen, Efficient Degradation of Toxic Organic Pollutants with Ni₂O₃/TiO₂-xBx under Visible Irradiation, Journal of the American Chemical Society 126, 2004, 4782-4783.
6. K.Torniaine, S.Tammilehto, V.Ulvi, The effect of pH, buffer type and drug concentration on the photodegradation of ciprofloxacin, International Journal of Pharmaceutics, 132 , 1996, 53–61.