



# INTERNATIONAL JOURNAL OF PURE AND APPLIED RESEARCH IN ENGINEERING AND TECHNOLOGY

A PATH FOR HORIZING YOUR INNOVATIVE WORK



## SPECIAL ISSUE FOR INTERNATIONAL LEVEL CONFERENCE "ADVANCES IN SCIENCE, TECHNOLOGY & MANAGEMENT" (IC-ASTM)

### A FACILE SYNTHESIS OF SOME NEW N- LACTOSYLATED THIOCARBAMATES NANOPARTICLES AND COMPARATIVE ACCOUNT OF THEIR ANTIMICROBIAL ACTIVITY WITH BULK SOLUTION

POONAM T. AGRAWAL

P.G. Department of Chemistry, Shri R.L.T. College of Science, Akola-444001 (M.S.)

Accepted Date: 05/09/2017; Published Date: 10/10/2017

**Abstract:** Nanoparticles are of great scientific interest as they are effectively a bridge between bulk materials and atomic or molecular structures. Nanoparticles research is currently an area of intense scientific research, due to a wide variety of potential application in biomedical, optical and electronic fields. In view of this it appeared quite interesting to prepared Nanoparticles of organic compounds containing carbohydrates

**Keywords:** Lactosyl thiocarbamides, Nanoparticles and Antimicrobial activity.



PAPER-QR CODE

Corresponding Author: POONAM T. AGRAWAL

Co Author: -

Access Online On:

[www.ijpret.com](http://www.ijpret.com)

How to Cite This Article:

Poonam T. Agrawal, IJPRET, 2017; Volume 6 (2): 315-318

## INTRODUCTION

In view of applications of lactosyl thiocarbamates and the nanoparticles in medicinal chemistry and in many other ways<sup>1</sup>, we herein report the synthesis of 1-hepta-o-benzoyl  $\beta$ -D-lactosyl-O-aryl - thiocarbamates nanoparticles by the use of ultrasonicator.

Nanoparticles are at the leading edge of the rapidly developing field of nanotechnology. Their unique size depended properties make these materials superior and indispensable in many areas of human activity. In recent years, nanoparticles are gaining importance due to their unique properties and their antimicrobial activities that are significantly different from those of bulk materials<sup>2</sup>.

Nanostructure materials are attracting a great deal of attention because of their potential for achieving specific processes and selectivity, especially in biological and pharmaceutical applications<sup>3</sup>. Recent studies have demonstrated that especially formulated nanoparticles have good antibacterial activity<sup>4,5</sup>.

### Experimental:

Specific rotations were measured on Equip-Tronics Digital Polarimeter at 28 °C in  $\text{CHCl}_3$ . IR spectra were recorded on Perkin-Elmer spectrum RXI FTIR spectrophotometer (4000-450  $\text{cm}^{-1}$ ).  $^1\text{H}$  NMR was recorded in  $\text{CDCl}_3$  on Bruker DRX-300 spectrometer operating at 300 MHz. The mass spectra were recorded on Jeol-SX-102(FAB) instrument.

#### a) Preparation of lactose octabenzoate:

In a 1 litre bottle having a tight cork, 55 ml dry pyridine and 55 ml of dry chloroform was taken. The bottle was cooled in an ice-salt bath. Now to this solution previously prepared cooled solution of 55 ml of benzoyl chloride in 55 ml dry chloroform was added with constant stirring. To this solution 20 gm of dry powder of lactose was added in several installments with constant stirring and maintaining the temperature of the reaction mixture below 5 °C. This solution was allowed to stand for 24 hr, it was then transferred to a 500 ml conical flask. The solution was washed several times with dilute aqueous sulphuric acid, water. The solution layer was separated by separating funnel. Afterwards the chloroform was removed, a white precipitate was isolated with petroleum ether and purified with chloroform ether with m.p.114 °C.

#### b) Preparation of brominating reagent:

Glacial acetic acid (30 ml) was taken in a conical flask and to it was added red phosphorous (3.0 gm). To this mixture molecular bromine (7 ml) was added gradually with constant shaking and cooling. The resultant mixture was allowed to stand at ice cold temperature for about a 30 min.

#### c) Synthesis of hepta-O-benzoyl- $\alpha$ -D-lactosyl bromide:

The finally powdered lactose octabenzoate (0.03M, 21.0g) was added gradually to the brominating agent. After the addition the flask was kept for 2hr at room temperature. Then the reaction mixture with chloroform (130ml) then the mixture was shaken vigorously for about 15 min. The resultant mixture was poured into ice cold water. The chloroform layer was then separated. It was washed several with aqueous sodium bicarbonate to remove excess of acetic acid followed by aqueous sodium metabisulphite to remove excess of bromine and finally 2-3 times with water. To the chloroform addition of petroleum ether afforded a solid (16.5 gm). This solid was expected hepta-O-benzoyl- $\alpha$ -D-lactosyl bromide (yield 77%). It was purified by dissolving it in minimum quantity of chloroform and reprecipitating it with petroleum ether m.p.168 °C.

#### (2)Preparation of lead thiocyanate :

Lead thiocyanate was prepared by mixing aqueous solution of lead nitrate and ammonium thiocyanate. The white granular lead thiocyanate was filtered washed with distilled water and dried at 50 °C.

#### (3)Preparation of hepta-O-benzoyl- $\beta$ -D-lactosyl isothiocyanate:

To a suspension of hepta-O-benzoyl- $\alpha$ -D-lactosyl bromide (21 gm, 0.03M) in sodium dried xylene (80ml) was added lead thiocyanate (6gm, 0.03M). The reaction mixture was then treated for microwave synthesis for about 3 min. This solution was then cooled and liberated lead bromide was removed by filtration. The xylene filtrate was then treated with petroleum ether (60-80 °C) with stirring, a white solid mass obtained (13gm). This solid was expected hepta-O-benzoyl-  $\beta$ -D-lactosyl isothiocyanate.

It was purified by dissolving it in minimum quantity of chloroform and reprecipitating it with petroleum ether, m.p. 118-120°C. [found; C;67.07, H;4.46, N;1.22, S;2.9; C<sub>62</sub>H<sub>49</sub>O<sub>17</sub>NS requires; C;66.96, H;4.41, N;1.26, S;2.88%].

**(4) Preparation of hepta-O-benzoyl-β-D-lactosyl -O-aryl thiocarbamates:**

To a suspension of lactosyl isothiocyanate aromatic alcohols were added in the reaction mixture was then treated in microwave for about 5 mins. This solution was then cooled and liberated lead bromide was removed by filtration. The xylene filtrate was then treated with petroleum ether (60-80°C) with stirring, a white solid mass obtained. This solid was expected hepta-O-benzoyl-β-D-lactosyl -O-aryl thiocarbamates.

**(4)Preparation of Nanoparticles of hepta-O-benzoyl-β-D-lactosyl -O-phenyl thiocarbamates:**

Take about 1 gm of hepta-O-benzoyl-β-D-lactosyl O-aryl thiocarbamates and dissolve it completely in the 50ml of solvent in 250 ml beaker. Now put this beaker in sonicator. The highly penetrating acoustic waves are passed through mixture, which create high pressure bubbles in the beaker due to which breakdown of the bulk material is takes place and desired sized nanoparticles are formed. The size determination of nanoparticles are done by the X-ray diffraction studies.

**IR SPECTRUM OF 1-HEPTA-O-BENZOYL-β-D-LACTOSYL -O-PHENYL THIOCARBAMATES<sup>6</sup>:-**

Absorption observed (Cm <sup>-1</sup> )	Assignment	Absorption Expected (Cm <sup>-1</sup> )
3066	C-H Ar Stretching	3040-3010
1729	C=O stretching	1750-1735
1176	C-O Stretching	1210-1153
1068,909	Charecteristic of Lactose	1100-1000 and 910-900

**NMR SPECTRAL STUDIES:** The NMR Spectrum <sup>7,8</sup> of compound distinctly displayed signals due, Aromatic Protons at δ 7.47-7.15 ppm, lactosyl protons at δ 5.77-3.76 ppm.

**Characterisation of Nanoparticles:**

**1.Characterisation using UV – Visible Spectrophotometer:** Characteirisation of nanoparticles was done using visible Spectrophotomter by using model Single Beam UV-Visible Spectrophotometer with software(BI/CI/SP/SB-S-03)of Bio Era make. The UV-Visible Spectroscopy reveals the formation of nanoparticles by showing different absorption those from bulk materials.

1. **Size determination of Lactose Thiocarbamates Nanoparticle by X-Ray Diffraction Studies:** From the X-Ray diffraction it comes to know that size of nanoparticles are as follows.

Compound		Size of Nanoparticles in nm
1-hepta-O-benzoyl-β-D-lactosyl thiocarbamates(6a)	-O- phenol	135
1-hepta-O-benzoyl-β-D-lactosyl thiocarbamates(6b)	- O-resorcinol	148
1-hepta-O-benzoyl-β-D-lactosyl thiocarbamates(6c)	-O- p-crysol	167

**ANTIMICROBIAL ACTIVITY COMPARISON:**

The bulk Lactosyl thiocarbamates and the Nanopartices of Lactosyl thiocarbamates have been screened for antibacterial activity using cup plate agar diffusion method by measuring the inhibition zone in mm. The compounds were taken at a concentration of 1 mg/ ml using dimethyl sulphoxide as solvent. Amikacin (100µg/ml ) was used as a standard for antibacterial activity. The compounds were screened for antibacterial activity against *Escherichia coli*, *Staphylococcus aureus*, in nutrient agar medium.

Antimicrobials	Zone of inhibition in mm					
	Bulk (5a)	Nanoparticles (6a)	Bulk (5b)	Nanoparticles (6b)	Bulk (5c)	Nanoparticles (6c)
<i>E. coli</i>	11	15	12	16	11	15
<i>S. aureus</i>	12	16	11	18	12	18
<i>S. typhi</i>	10	15	12	15	10	16
<i>P. vulgaris</i>	10	17	12	14	10	14
Amikacin	11	20	10	18	12	18
Clandamycine	11	18	11	15	10	15

\*including the well diameter of 8mm. \*\* zone of inhibition in mm (15or less) resistance, (16- 20mm) moderate and (more than 20mm) sensitive

**Conclusion:** Lactosyl thiocarbamates Nanoparticles show good antimicrobial activity as compare to the bulk solution of lactosyl thiocarbamates due to their large surface area to volume ratio, which is coming up current interest in the researchers.

#### ACKNOWLEDGEMENT:

Authors are thankful to RSIC, CDRI Lucknow for providing the spectra, to Dept. of Biotechnology, North Maharashtra University, Jalgaon for providing X-Ray Diffractions.and also to Dr. S.P.Deshmukh, Shri Shivaji College, Akola, Dr. V.D.Nanoty, Principal Shri R.L.T. College of Science, Akola for providing Guidance and necessary facilities.

#### REFERENCES:

1. C. H. Cao, C.J.Zhou, H.Y.Gao, Y.T. Liu: *J.Chin .Chem.Soc.*, 2001, **48**,207-210
2. L.Mazzola,Nature Biotechnology .,2003.,**21**,1137-1147.
3. M. Maillard, S.Giorgo.,M.P.Pilent *Adv. Material.*,2002.,**14(15)**,1084-1086.
4. Z.Chen.,L.Gao*Material Research Bulletin* .,2007.,**42**,1657-1661.
5. L. Mazur, *Electrochemistry Communication.*,2004.,**6**,400-403.
6. R.M. Silverstein, G.C. Bassler and T. C. Morrill. : "Spectrometric Identification of Organic Compounds," 5th Ed.;John wiley and sons, INC, New York, 2003,P 108, 119, 120, 123.
7. N.B.Colthup, L.H.Daly and S.E.Wiberley. : *Introduction to Infrared and Raman Spectroscopy*," Academic Press, New York, 2003, P-279.
8. D.H.williams and I.Flemming "Spectroscopic methods in organic chemistry" 4 th Ed., Tata McGraw-Hill Publication New Delhi, 2003, P 40, 41, 47, 53.